

- Semi-intensive farming
- High performance
- High survival
- Medium energy starter diet



COMPOSITION:

Analyses (%)	0.2-0.3 mm	0.3-0.5 mm	0.5-0.8 mm	1.00 mm	1.5 mm
Protein	56	56	56	54	54
Fat	15	15	15	15	15
Crude fibre	0,3	0,3	0,3	0,3	0,3
Ash	11,30	11,30	11,30	11,96	11,96
Total P	1,77	1,77	1,77	1,75	1,75

Vitamins added

	0.2-0.3 mm	0.3-0.5 mm	0.5-0.8 mm	1.00 mm	1.5 mm
Vitamin A (IE/kg)	16085	16085	16085	12000	12000

Energy (MJ/kg)

	0.2-0.3 mm	0.3-0.5 mm	0.5-0.8 mm	1.00 mm	1.5 mm
Gross Energy	21,2	21,2	21,2	20,6	20,6
Digestible Energy	19,2	19,2	19,2	19,0	19,0

FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	> 24 °C
< 0.2	0.2-0.3		2,66	3,22	3,89	4,71	5,69	6,89	5,58	
0.2-0.5	0.3-0.5	According to fish's appetite	2,44	2,95	3,57	4,32	5,22	6,32	5,12	According to fish's appetite & O2 level
0.5-1.5	0.5-0.8		2,18	2,64	3,19	3,86	4,67	5,65	4,58	
1.5-5.0	1.0		1,66	2,00	2,42	2,93	3,54	4,29	3,47	
5.0-10	1.0/1.5		1,33	1,61	1,95	2,35	2,85	3,44	2,79	
10-30	1.5		0,96	1,17	1,41	1,71	2,06	2,49	2,02	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.