

- Semi-intensive farming
- Good performance
- Very palatable
- Optical feeding control



COMPOSITION:

Analyses (%)

Analyses (%)		Sizes
Protein	42	3.0 mm
Fat	13	4.5 mm
Crude fibre	2,7	6.0 mm
Ash	13,0	
Total P	1,39	

Vitamins added

Vitamin A (IE/kg)	10002
-------------------	-------

Energy (MJ/kg)

Gross Energy	19,4
Digestible Energy	15,8

FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 10 °C	10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	26 °C
50-100	3,0		0,85	1,03	1,25	1,51	1,82	2,21	2,67	3,23	2,90	
100-250	4,5	According to fish's appetite	0,66	0,80	0,97	1,17	1,42	1,72	2,08	2,51	2,26	According to fish's appetite & O2 level
250-500	6,0		0,57	0,69	0,83	1,01	1,22	1,47	1,78	2,16	1,94	
500-1000	6,0		0,50	0,60	0,73	0,88	1,06	1,28	1,55	1,88	1,69	
1000-2000	6,0		0,43	0,52	0,63	0,76	0,92	1,12	1,35	1,63	1,47	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.