

- For semi-intensive farming
- Good performance



## COMPOSITION:

### Analyses (%)

|             |      |       |        |
|-------------|------|-------|--------|
| Protein     | 45   | Sizes | 2.0 mm |
| Fat         | 18   |       |        |
| Crude fibre | 1,3  |       |        |
| Ash         | 9,3  |       |        |
| Total P     | 1,50 |       |        |

### Vitamins added

|                   |       |
|-------------------|-------|
| Vitamin A (IE/kg) | 11877 |
|-------------------|-------|

### Energy (MJ/kg)

|                   |      |
|-------------------|------|
| Gross Energy      | 21,2 |
| Digestible Energy | 18,9 |

## FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


| Fish weight (g) | Feed size (mm) | < 12 °C                      | 12 °C | 14 °C | 16 °C | 18 °C | 20 °C | 22 °C | 24 °C | > 24 °C                                 |
|-----------------|----------------|------------------------------|-------|-------|-------|-------|-------|-------|-------|---|
| 30-50           | 2.0            | According to fish's appetite | 0,84  | 1,01  | 1,22  | 1,48  | 1,79  | 2,16  | 1,75  | According to fish's appetite & O2 level |
| 50-100          | 2.0            |                              | 0,75  | 0,91  | 1,10  | 1,33  | 1,61  | 1,95  | 1,58  |   |

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

## ECOLOGICAL FIGURES:

Discharge per 1000 kg production



### Feed conversion ratio

The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.