



- Broodstock diet
- Optimal egg development
- High egg quality and fry survival

COMPOSITION:

Analyses (%)		Sizes
Protein	48	6.0 mm
Fat	15	9.0 mm
Crude fibre	1,2	
Ash	8,9	
Total P	1,36	
Astaxanthin (mg/kg)	40	

Vitamins added

Vitamin A (IE/kg)	24998
-------------------	-------

Energy (MJ/kg)

Gross Energy	20,7
Digestible Energy	18,5

FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	> 24 °C
1000-2000	6.0		0,32	0,43	0,57	0,69	0,84	1,01	0,80	
2000-4000	6.0	According to fish's appetite	0,23	0,30	0,40	0,48	0,59	0,71	0,56	According to fish's appetite & O2 level
4000-6000	6.0		0,18	0,24	0,32	0,39	0,47	0,57	0,45	
6000-8000	6.0-9.0		0,15	0,19	0,26	0,31	0,38	0,46	0,36	
8000-15000	9.0		0,09	0,11	0,15	0,18	0,22	0,27	0,21	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.